

COMMUNITY USER WORKSHOP ON PLANETARY LIBS (CHEMCAM) DATA

Introduction to LIBS



Sam Clegg

Los Alamos National Laboratory

sclegg@lanl.gov



Creating LIBS Sparks



← 3" →



LIBS Spots and Dust Removal

ChemCam
Dust Removal


ChemCam laser marks

Brushed
Surface

“Wernecke”
Sol 169




NASA/JPL-Caltech/MSSS/Honeybee
Robotics/LANL/CNES

NASA/JPL-
Caltech/LANL/IRAP/CNES/LPGNantes/I
AS/CNRS/MSSS






Ca LIBS Spectrum-LANL Testbed



- Not every NIST emission line is observed in a LIBS spectrum.
 - http://physics.nist.gov/PhysRefData/ASD/lines_form.html
 - Every LIBS line must be found in NIST
 - Typically observe neutral and first ionized atoms, perhaps second ionization under ChemCam conditions.
- Must be Spectrally well calibrated!
 - Know the difference between vacuum vs. air (Earth or Mars) calibration
 - Closest NIST emission line is not good enough
- More details provided in the next few presentations




Raw ChemCam Spectrum – Jake_1






Raw ChemCam Spectrum – Jake_1




Raw ChemCam Spectrum – Jake_1



Carefully Processed Spectra Lead to Quantitative Analysis
Continuum Removal, Spectral Calibration, Distance Correction are Critical


ChemCam Spectra



Fully Processed Spectra Ready for Quantitative Analysis
 Much can be Qualitatively Observed

Chemical Matrix Effects Complicate Quantitative Analysis

Peak Area Analysis Method



- Conventional Elemental Analysis
 - Peak Area or Height vs. Concentration
 - Each Peak is Analyzed Independently
- Sample Elemental and Molecular Composition Influences:
 - Laser-to-Sample Coupling Efficiency
 - Chemical Reactions within the Plasma
 - Collisional Quenching
- Chemical Matrix Effects
 - Increase Scatter and Uncertainty
- Chemical Matrix Effects Compensation
 - Cal-Free LIBS
 - Various Normalization

Multivariate analyses are used to compensate for these matrix effects

Quantitative Calibration

3 m standoff distance			
BHVO-2	DH 4912	Norite	Swy-2
GBW 07105	JR-1	GYP A	SGR-1
NBS688	GBW 07113	GYP C	VS MO7
BIR-1	Ultramafic*	GYP D	UNS ZK
BCR-2	Umph*	MHC1356*	GUW GNA
JA-1	Cadillac*	MHC2319*	M6 Haggerty*
Ja2	VH-1*	VZO106	GYP B
Ja3	MSHA*	VZO114	MHC3828*
MO12	Moppin*	NAu2-Hi-S	UNS AK
MO14	BK-2	NAu2-Med-S	GBW 07313
JB-2	BWQC1*	NAu2-Lo-S	GBW 07316
GSR-2	Trond*	KGa2-Med-S	SARM51
BE-N	WMG*	NBS-88b	STSD-1
AGV-2	VH-49*	JDo-1	STSD-3
JB-3	Grano Dike*	GBW 07108	STSD-4
BT-2	Macusanite	NBS 97a	
GBW 07110	Picrite	NBS 98a	
GBW07104	Shergottite	NAu2	


	Igneous
	Sedimentary
	Sulfur-rich
	Rover cal

* = from Dyar lab, all others from commercial sources

backup



Ca Electronic Energy Level Diagram



Emission lines are produced as atoms relax from upper state to lower state.
 Lower state is not always the ground state.